Do platelet mediators and coagulation factors affect the formation of neutrophil extracellular traps?

Iraklis C. Moschonas, Styliani Papadaki, Sofia Sidiropoulou, Alexandros D. Tselepis

Atherothrombosis Research Centre/Laboratory of Biochemistry, Department of Chemistry,
University of Ioannina, Ioannina, Greece

ABSTRACT

Aim: To investigate the effect of factor Xa (FXa), thrombin and platelets, under activation or resting conditions, on neutrophil extracellular traps (NETs) formation (NETosis). **Materials and Methods:** Neutrophils, isolated from apparently healthy volunteers, were challenged with 25-50 nM FXa, 0.2-8 U/mL thrombin, 5-100 μ M adenosine diphosphate (ADP), 0.1-0.5 mM arachidonic acid (AA), 5-15 μ g/mL collagen, 100 μ M protease-activated receptor (PAR)-4 activating peptide, 10-100 μ M PAR-2 activating peptide or 25-100 nM phorbol 12-myristate 13-acetate (PMA). Moreover, neutrophils were challenged with platelet-rich plasma (PRP), activated with ADP (at a concentration of either 20 μ M, in an aggregometer, or of 50 μ M, *in situ*). Neutrophils were incubated for 3.5 h at 37°C and 5% CO₂ and an immunofluorescence protocol followed, in order to stain neutrophils and NETs for myeloperoxidase and DNA. Finally, NETs and neutrophils were microscopically visualized and the percentage of NET-releasing neutrophils of each of the above experimental conditions was evaluated. **Results and Conclusion:** Among the agonists studied, only PMA and AA, induced robust NETosis, at a threshold concentration of 25 nM and 0.25 mM, respectively. Furthermore, platelets pre-activated with ADP, but not resting platelets induce the formation of NETs, a phenomenon that is also observed when platelets were activated with ADP *in situ*, in the presence of neutrophils. The present results provide an additional mechanism through which platelets contribute to various pathophysiological conditions in which NETs are implicated, such as inflammation and thrombosis. The whole spectrum of platelet mediators that are implicated in NETosis needs to be further investigated.

KEY WORDS: Arachidonic acid, factor Xa, neutrophil extracellular traps, platelets, thrombin

1. INTRODUCTION

Neutrophil extracellular traps (NETs) are threads of de-condensed chromatin, which are decorated with numerous

proteins and are generated by activated neutrophils.¹ The procedure of the formation of NETs is termed as NETosis. Originally, NETs were studied for their anti-microbial properties, since they entrap and possibly kill pathogens, by providing a high local concentration of microbicidal proteases, such as neutrophil elastase and myeloperoxidase (MPO).¹ However, later, extended research, has revealed that NETs are also present under sterile inflammatory

Corresponding author:

Alexandros D. Tselepis, MD, PhD, Professor of Biochemistry-Clinical Chemistry, Atherothrombosis Research Centre/Laboratory of Biochemistry, Department of Chemistry, University of Ioannina, 45110, Ioannina, Greece
Tel.: +30 26510 08365, Fax: +30 26510 8785
E-mail: atselep@uoi.gr

Submission: 05.06.2020, Acceptance: 15.06.2020

conditions and they participate in the pathogenesis of multiple diseases, e.g. cancer,² diabetes mellitus³ and Alzheimer's disease.⁴ Importantly, it is established that NETs are heavily involved in atherogenesis, as well as in arterial and venous thrombosis.⁵

In addition to their central role in the coagulation cascade, coagulation factors Xa (FXa) and thrombin exert cellular effects via protease-activated receptors (PARs),^{6,7} which are expressed on virtually all cell types. FXa activates PAR-1 and -2, whereas thrombin activates PAR-1, -3 and -4, but not PAR-2.^{6,7}

Platelets play a key role in hemostasis and thrombosis, however they also interact and activate leukocytes, including neutrophils, thus participating in inflammatory responses as well as in atherogenesis.⁸ The platelet-neutrophil interaction is mediated by P-selectin, expressed on the platelet surface, and P-selectin glycoprotein ligand-1, which is expressed on the surface of neutrophils.⁸

In the present study, we investigated the effect of FXa and thrombin, as well as platelets, in their resting and activation conditions, on neutrophils activation and NETs production.

2. MATERIALS AND METHODS

2.1. Reagents

Adenosine diphosphate (ADP), arachidonic acid (AA), thrombin from human plasma, Histopaque®-1119, heat inactivated (HI) human serum from clotted whole blood, phorbol 12-myristate 13-acetate (PMA) and 4',6-diamidino-2-phenylindole (DAPI) were purchased from Sigma-Aldrich (St. Louis, MO, USA). Collagen type IV was purchased from Chrono-Log Corporation (Philadelphia, PA, USA). PAR-2 and -4 activating peptides were purchased from Bachem (Basel, Switzerland). FXa was purchased from Haematologic Technologies, Inc. (Essex Junction, VT, USA). Anti-MPO antibody was purchased from Santa Cruz (Dallas, TX, USA). Ficoll® Paque Plus was purchased from GE Healthcare (Chicago, IL, USA). Roswell Park Memorial Institute (RPMI) Media 1640 was purchased from Gibco (Thermo Fisher Scientific, Waltham, MA, USA). BioCoat Cellware poly-L-lysine 12 mm coverslips were purchased from Becton Dickinson (La Jolla, CA, USA). Goat anti-rabbit IgG (H+L) secondary antibody, Alexa Fluor® 488, was purchased from Invitrogen (Thermo Fisher Scientific, Waltham, MA, USA).

2.2. Isolation of neutrophils

Heparinized whole blood was withdrawn from fasting, non-smoking, apparently healthy volunteers, who had not received any anti-platelet and anti-inflammatory drugs for at least two weeks prior to venipuncture. The whole blood was diluted with normal saline, it was care-

fully overlaid on a Histopaque®-1119 and Ficoll® Paque Plus gradient and then it was centrifuged at $700 \times g$ for 30 min, at room temperature (RT). The polymorphonuclear leukocyte (PMN) zone was aspirated, diluted with 12.5 mM phosphate-buffered saline (PBS) and centrifuged at $200 \times g$ for 10 min, at RT. Finally, the PMN pellet, enriched in neutrophils, was re-suspended with RPMI Media 1640, supplemented with 2% HI human serum.

2.3. Preparation of platelet-rich plasma

Platelet-rich plasma (PRP), autologous to the neutrophils, was prepared as previously described.⁹ Briefly, whole blood was centrifuged at $126 \times g$ for 15 min, at RT. The upper two-thirds of the PRP layer were aspirated and the remaining PRP was further centrifuged at $1500 \times g$ for 20 min, at RT, to obtain platelet-poor plasma (PPP). The number of platelets in the PRP was counted using a hemocytometer and it was adjusted to $2.5 \times 10^5/\mu\text{L}$ with PPP.

2.4. NETs generation and visualization

Neutrophils (2×10^5) were added to 24-well plates, in which poly-L-lysine coverslips were pre-placed, and incubated for 1 h at 37°C and 5% CO₂, in order to allow the cells to bind to the coverslips.

The effect of coagulation factors and various classic platelet agonists on NETs generation was investigated. We tested FXa (25-50 nM), thrombin (0.2-8 U/mL), ADP (5-100 μM), AA (0.1-0.5 mM), collagen (5-15 $\mu\text{g/mL}$) and PAR-4 activating peptide (sequence AYPGKF-NH₂; 100 μM). In addition, we studied the effect of a selective, PAR-2 activating peptide (sequence SLIGRL-NH₂; 10-100 μM) on the formation of NETs, since PAR-2 is expressed on neutrophils.¹⁰ PMA, a widely used NET inducer, was used as a positive control for induction of NETosis at a concentration range of 25-100 nM.

In other experiments, PRP was activated with 20 μM ADP at 37°C for 5 min, under 1200 RPM stirring conditions, in a Chrono-Log Model 700 Lumi-Aggregometer (Chrono-Log Corporation, Philadelphia, PA, USA) and the total suspension of the activated PRP was then immediately transferred to neutrophils, at a physiological ratio of 150×10^3 platelets/ 6×10^3 neutrophils. Alternatively, PRP was added to neutrophils, at the above ratio, and activated *in situ* with 50 μM ADP.

In all the above experimental conditions, neutrophils were incubated for 3.5 h at 37°C and 5% CO₂, to produce NETs. Afterwards, the cells and NETs were fixed and then an immunofluorescence protocol was followed. According to this protocol, neutrophils and NETs were incubated with an anti-MPO primary antibody and the goat anti-rabbit IgG (H+L) secondary antibody, Alexa Fluor® 488, for MPO

visualization. In addition, neutrophils and NETs were incubated with 200 ng/mL DAPI, for DNA visualization. NETs were imaged with an Olympus BX41 fluorescence microscope and with a Leica TCS SP5 confocal microscope, with a HC PL APO CS 63× oil lens and the LAS AF software. The number of cells that release NETs was counted, and the percentage of NET-releasing neutrophils was obtained from 5 representative optical fields.

3. RESULTS AND DISCUSSION

The coagulation factors FXa and thrombin were tested at a range of concentrations of 25-50 nM and 0.2-8 U/mL, respectively. Even at the maximal concentration tested, FXa or thrombin did not induce substantial NETosis (Figure 1).

To study the effect of activated platelets and therefore the effect of platelet - neutrophil cell to cell interaction or the effect of platelet-secreted mediators, we first studied the effect of various platelet agonists on NETosis, in a wide variety of concentrations. Among all agonists tested, only PMA and AA induced robust NETosis (Figure 2A). The threshold concentration of the above agonists that induced the maximum of NETs production, were 25 nM for PMA and 0.25 mM for AA. All other agonists, i.e. ADP, collagen, PAR-4 and PAR-2 agonists, even at the maximum concentration tested, did not induce substantial NETosis (Figure 2A). Representative confocal microscopy images of untreated neutrophils, as well as PMA- and AA-induced NETosis, at the maximum concentration tested, are shown in Figure 2B.

We next evaluated whether activated platelets can induce NETosis. To activate platelets *in situ*, i.e. in the presence of neutrophils, we have chosen a platelet agonist, among those tested in the present study, that was inactive towards NETs production by neutrophils. Among these

FIGURE 1. Effect of coagulation factors on the formation of NETs. The bar graph depicts NETosis induced by 50 nM FXa and 8 U/mL thrombin. The above results represent the mean $\pm$ SD of at least 4 different neutrophil preparations.

FIGURE 2. A. Effect of various agonists on the formation of NETs. The bar graph depicts NETosis induced by 25 nM PMA, 100 μ M ADP, 0.25 mM AA, 15 μ g/mL collagen, 100 μ M AYPGKF-NH₂ and 100 μ M SLIGRL-NH₂. The above results represent the mean $\pm$ SD of at least 4 different neutrophil preparations. **B.** Representative confocal microscopy images of untreated neutrophils, as well as NETs from neutrophils treated with 25 nM PMA or 0.25 mM AA. Neutrophils and NETs were stained for myeloperoxidase (MPO) and DNA with DAPI. Magnification 63×. Abbreviation: N; neutrophils.

agonists we have chosen ADP since its platelet receptor P2Y₁₂, is the main target of antiplatelet therapy used currently in the clinical practice.¹¹ ADP cannot induce neutrophil activation and NETosis, since neutrophils do not express functional P2Y₁₂ receptor.¹² We first investigated whether the total suspension of platelets activated with ADP in an aggregometer could induce NETosis. As expected, ADP, at the concentration of 20 μ M, induced a full platelet aggregatory response when added to the

PRP in the aggregometer (Figure 3A). The total suspension of the ADP-activated PRP induced a significant NETosis when incubated with the neutrophil suspension for 3.5 h as compared with resting PRP (Figure 3B). Similarly, when the PRP was activated with ADP *in situ*, in the presence of neutrophils, it induced NETs production, which was not observed in the absence of ADP (Figure 3C). Representative confocal microscopy images of neutrophils in the presence of untreated PRP, as well as in the presence of PRP, either pre-activated in the aggregometer or activated *in situ* with ADP, are shown in Figure 3D.

The present study demonstrates that the coagulation factors FXa and thrombin, as well as the platelet agonists ADP, collagen and PAR-4 agonist, do not induce NETs production. This is possibly due to the absence of the expression of their corresponding functional receptors on neutrophils, or due to the fact that these receptors may be

linked to other neutrophil functions, such as chemotaxis, but not to NET formation.¹²⁻¹⁴ It has been suggested that PAR-2 is expressed in neutrophils.¹⁰ However, in the present study we showed that the PAR-2 activating peptide does not cause substantial NETosis. This could be possibly due to the fact that this receptor may be linked to other neutrophil functions but not to NET formation, or this receptor may be not functional in neutrophils.

By contrast the platelet agonist AA, induced robust NETosis, comparable in magnitude to PMA, the potent neutrophil activator and inducer of NETs production. In platelets, AA is physiologically hydrolyzed from the membrane bilayer phospholipids, or taken up by them when added *in vitro*, to form thromboxane A₂ via the cyclooxygenase-1 pathway.¹⁵ In neutrophils, AA is metabolized via the lipoxygenase-1 pathway, which among other active metabolites, it leads to the formation of leukotriene B₄,¹⁶

FIGURE 3. A. Representative aggregation curve of the response of PRP, activated with 20 μ M ADP. **B.** Effect of PRP, treated with 20 μ M ADP in an aggregometer, on the formation of NETs (* $p = 0.03$, compared with either PPP or untreated [resting; Rest.] PRP). **C.** Effect of PRP, treated with 50 μ M ADP *in situ*, on the formation of NETs (** $p = 0.011$, compared with either PPP or untreated [resting; Rest.] PRP). **D.** Representative confocal microscopy images of neutrophils in the presence of untreated PRP, as well as in the presence of PRP, either pre-activated in the aggregometer or activated *in situ* with ADP. Neutrophils and NETs were stained for myeloperoxidase (MPO) and DNA with DAPI. Magnification 63 $\times$. Abbreviation: N; neutrophils.

that could be the main metabolite responsible for the AA-induced NETosis.

An important observation of the present study is that in the presence of platelets, ADP can induce NETosis even though it is inactive when added directly to neutrophils. This finding is consistent with our observation that the suspension of platelets activated with ADP, also induces NETosis. These results suggest that activated platelets, through cell to cell interaction with neutrophils or by secreting mediators such as P-selectin or AA or by shedding platelet microparticles could potentially induce the production of NETs.

In conclusion, the present study demonstrates that activated platelets induce NETs production thus providing an additional mechanism through which platelets contribute to various pathophysiological conditions in which NETs are implicated, such as inflammation and

thrombosis. This new finding may be clinically significant in patients receiving antiplatelet therapy with the platelet P2Y₁₂ antagonists clopidogrel or ticagrelor. The whole spectrum of platelet mediators that are implicated in NETosis needs to be further investigated.

Acknowledgments

The present research has been co-financed by the Operational Program "Human Resources Development, Education and Lifelong Learning" and is co-financed by the European Union (European Social Fund) and Greek national funds [grant number (MIS) 5006031].

Conflict of interest

The authors report no conflicts of interest.

ΠΕΡΙΛΗΨΗ

Επηρεάζουν οι αιμοπεταλιακοί διαμεσολαβητές και οι παράγοντες πήξης το σχηματισμό ουδετεροφιλικών δικτύων χρωματίνης (NETs);

Ηρακλής Χ. Μοσχονάς, Στυλιανή Παπαδάκη, Σοφία Σιδηροπούλου, Αλέξανδρος Δ. Τσελέπης

Ερευνητικό Κέντρο Αθηροθρόμβωσης/Εργαστήριο Βιοχημείας, Τμήμα Χημείας, Πανεπιστήμιο Ιωαννίνων, Ιωάννινα, Ελλάδα

Σκοπός: Η διερεύνηση της δράσης των παραγόντων πήξης Χα (factor Χα, FΧα), θρομβίνης και των αιμοπεταλίων, υπό συνθήκες ενεργοποίησης ή εν ηρεμία, στο σχηματισμό ουδετεροφιλικών δικτύων χρωματίνης (neutrophil extracellular traps, NETs· νέτωση). **Υλικά και Μέθοδοι:** Ουδετερόφιλα, απομονωμένα από φαινομενικά υγιείς εθελοντές, επιδράσθηκαν με 25-50 nM FΧα, 0,2-8 U/mL θρομβίνης, 5-100 μM διφωσφορικής αδενοσίνης (adenosine diphosphate, ADP), 0,1-0,5 mM αραχιδονικού οξέος (arachidonic acid, AA), 5-15 μg/mL κολλαγόνου, 100 μM ενεργοποιητικού πεπτιδίου του υποδοχέα PAR-4, 10-100 μM ενεργοποιητικού πεπτιδίου του PAR-2 ή 25-100 nM οξικής μυριστικής φορβόλης (phorbol 12-myristate 13-acetate, PMA). Επιπλέον, τα ουδετερόφιλα επιδράσθηκαν με πλάσμα πλούσιο σε αιμοπετάλια (platelet-rich plasma, PRP), ενεργοποιημένο με ADP (συγκέντρωσης 20 μM σε συσσωρευόμετρο, ή 50 μM, *in situ*). Τα ουδετερόφιλα επώασθηκαν για 3,5 h στους 37°C και 5% CO₂ και ακολούθησε ανοσοφθορισμός με χρώση της μυελοϋπεροξειδάσης και του DNA. Τα ουδετερόφιλα και τα NETs παρατηρήθηκαν μικροσκοπικά και υπολογίστηκε το ποσοστό των ουδετεροφίλων που υπέστησαν νέτωση για κάθε μία από τις παραπάνω συνθήκες. **Αποτελέσματα και Συμπέρασμα:** Μεταξύ των αγωνιστών που μελετήθηκαν, μόνο το PMA και το AA, προκάλεσαν ισχυρή νέτωση, με συγκέντρωση κατωφλίου τα 25 nM και 0,25 mM, αντίστοιχα. Επιπλέον, αιμοπετάλια προενεργοποιημένα με ADP, αλλά όχι αιμοπετάλια εν ηρεμία, προκαλούν το σχηματισμό των NETs, ένα φαινόμενο που επίσης παρατηρείται όταν τα αιμοπετάλια ενεργοποιήθηκαν με ADP *in situ*, παρουσία ουδετεροφίλων. Τα αποτελέσματα αυτά παρέχουν έναν επιπρόσθετο μηχανισμό διαμέσου του οποίου τα αιμοπετάλια συνεισφέρουν σε διάφορες παθοφυσιολογικές καταστάσεις στις οποίες εμπλέκονται τα NETs, όπως είναι η φλεγμονή και η θρόμβωση. Το πλήρες φάσμα των αιμοπεταλιακών διαμεσολαβητών που εμπλέκονται στη νέτωση χρήζει περαιτέρω διερεύνησης.

ΛΕΞΕΙΣ ΚΛΕΙΔΙΑ: Αραχιδονικό οξύ, παράγοντας πήξης Χα, ουδετεροφιλικά δίκτυα χρωματίνης, αιμοπετάλια, θρομβίνη

REFERENCES

1. Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, et al., Neutrophil extracellular traps kill bacteria. *Science*. 2004 Mar;303(5663):1532-5.
2. Cools-Lartigue J, Spicer J, Najmeh S, Ferri L. Neutrophil extracellular traps in cancer progression. *Cell Mol Life Sci*. 2014 Nov;71(21):4179-94.
3. Wong SL, Demers M, Martinod K, Gallant M, Wang Y, Goldfine AB, et al, Diabetes primes neutrophils to undergo NETosis, which impairs wound healing. *Nat Med*. 2015 Jul;21(7):815-9.
4. Pietronigro EC, Della Bianca V, Zenaro E, Constantin G. NETosis in Alzheimer's Disease. *Front Immunol*. 2017 Mar;8:211.
5. Moschonas IC, AD Tselepis. The pathway of neutrophil extracellular traps towards atherosclerosis and thrombosis. *Atherosclerosis*. 2019 Sep;288:9-16.
6. Moschonas IC, Goudevenos JA, Tselepis AD. Protease-activated receptor-1 antagonists in long-term antiplatelet therapy. Current state of evidence and future perspectives. *Int J Cardiol*. 2015 Apr;185:9-18.
7. Papadaki S, Tselepis AD. Nonhemostatic Activities of Factor Xa: Are There Pleiotropic Effects of Anti-FXa Direct Oral Anticoagulants? *Angiology*. 2019 Nov;70(10): 896-907.
8. Kalantzi KI, Dimitriou AA, Milionis HJ, Goudevenos IA, Tselepis AD. Clopidogrel differentially affects platelet-mediated thrombosis and inflammatory response in patients with acute coronary syndromes. *J Thromb Haemost*. 2011 Apr;9(4):875-8.
9. Tsoumani ME, Tatsidou PT, Ntalas IV, Goudevenos JA, Tselepis AD. Dynamic platelet adhesion in patients with an acute coronary syndrome: The effect of antiplatelet therapy. *Platelets*. 2016 Dec;27(8):812-20.
10. Bock A, Tucker N, Kelher MR, Khan SY, Gonzalez E, Wohlauser M, et al. Alpha-Enolase Causes Proinflammatory Activation of Pulmonary Microvascular Endothelial Cells and Primes Neutrophils Through Plasmin Activation of Protease-Activated Receptor 2. *Shock*. 2015 Aug;44(2):137-42.
11. Tsoumani ME, Tselepis AD. Antiplatelet Agents and Anticoagulants: From Pharmacology to Clinical Practice. *Curr Pharm Des*. 2017;23(9):1279-93.
12. Le Duc D, Schulz A, Lede V, Schulze A, Thor D, Brüser A, et al. P2Y Receptors in Immune Response and Inflammation. *Adv Immunol*. 2017;136:85-121.
13. Kahn ML, Nakanishi-Matsui M, Shapiro MJ, Ishihara H, Coughlin SR. Protease-activated receptors 1 and 4 mediate activation of human platelets by thrombin. *J Clin Invest*. 1999 Mar;103(6):879-87.
14. Senior RM, Hinek A, Griffin GL, Pipoly DJ, Crouch EC, Mecham RP. Neutrophils show chemotaxis to type IV collagen and its 7S domain and contain a 67 kD type IV collagen binding protein with lectin properties. *Am J Respir Cell Mol Biol*. 1989 Dec;1(6):479-87.
15. Patrono C, Rocca B. Measurement of Thromboxane Biosynthesis in Health and Disease. *Front Pharmacol*. 2019 Oct;10:1244.
16. Raulf M, König W. Modulation of leukotriene generation from human polymorphonuclear granulocytes by polychlorinated biphenyls (PCB). *Immunology*. 1991 Aug;73(4):485-90.